

### Acetaminophen

Do not exceed 5 doses in 24 hours	May Give Every	10-11 lbs. 2-3 mo.	12-17 lbs. 4-11 mo.	18-23 lbs. 12-23 mo.	24-35 lbs. 2-3 yr.	36-47 lbs. 4-5 yr.	48-59 lbs. 6-8 yr.	60-85 lbs. 9-11 yr.
<u>Chewable 80 mg tablets</u>	4 - 6 hours				2 tab.	3 tab.	4 tab.	6 tab.
<u>Chewable Junior 160 mg</u>	4 - 6 hours						2 tab.	3 tab.
<u>DROPS 80 mg/0.8 ml</u> *	4 - 6 hours	1/2 dppr. 0.4 ml.	1 dppr. 0.8 ml.	1 1/2 dppr. 1.2 ml.	2 dppr. 1.6 ml.			
<u>Elixir 160 mg/5 ml</u>	4 - 6 hours		1/2 tsp.	3/4 tsp.	1 tsp.	1 1/2 tsp.	2 tsp.	3 tsp.
<u>Suppository 120 mg</u>	4 - 6 hours			1 supp.	1-1/2 supp.			
<u>Suppository 325 mg</u>	4 - 6 hours				1/2 supp.	3/4 supp.	1 supp.	
<u>Suppository 80 mg</u>	4 - 6 hours		1 supp.	1-1/2 supp.	2 supp.			

### Ibuprofen

Give every 6 hours for fever over 102° rectal	May Give Every	12-17 lbs. 6-11 mo.	18-23 lbs. 12-23 mo.	24-35 lbs. 2-3 yr.	36-47 lbs. 4-5 yr.	48-59 lbs. 6-8 yr.	60-85 lbs. 9-11 yr.
<u>Chewable Tablets 100 mg</u>	6 hours			1 tab.	1-1/2 tab.	2 tab.	2-1/2 tab.
<u>Chewable Tablets 50 mg</u>	6 hours			2 tab.	3 tab.	4 tab.	5 tab.
<u>DROPS 50 mg/1.25 ml</u> *	6 hours	1 dppr. 1.25 ml.	1-1/2 dppr. 1.875 ml.	2 dppr. 2.5 ml.	3 dppr. 3.75 ml.		
<u>Suspension 100 mg/5 ml</u>	6 hours	1/2 tsp.	3/4 tsp.	1 tsp.	1-1/2 tsp.	2 tsp.	2-1/2 tsp.
<u>Tablets 200 mg</u>	6 hours					1 tab.	1-1/2 tab.

\* **CAUTION:** We recommend that you use the dropper or dosage cup that comes with your medicine to avoid overdose. Dosage recommendations on bottles may vary from the recommendation on these pages. We would like you to use our recommended dosages unless your health care provider has given you a different (usually higher!) dosage. Aspirin should not be given to your child.